

PESCIENCE

Norcodrene™

Breakthrough Thermogenic†



▶ Thermo Igniting Matrix†

▶ Energy & Mood†

▶ Aframomum Extract

▶ **ChromaX**
chromium picolinate

Dietary Supplement

90
Capsules

Supplement Facts

Serving Size: 1 Capsule

Servings Per Container: 90

	Amount Per Serving	%DV
Chromium (as chromium picolinate) (Chromax®)	100 mcg	83%
Norcodrene Proprietary Blend	425 mg	*
Lemon Balm (<i>Melissa officinalis</i>) (leaf) Extract, L-Theanine, Aframomum Melegueta (seed) Extract, Kola Nut Extract		
Caffeine Anhydrous	100 mg	*

* Daily Value (DV) Not Established

Other Ingredients: Gelatin, Magnesium Stearate, Silicon Dioxide, Titanium Dioxide, FD&C Yellow #6, FD&C Green #3

Manufactured for PEScience LLC (3665 East Bay Dr #204-155, Largo, FL 33771, USA) Ph: 888-885-0195.

Chromax® and the Chromax logo are registered trademarks of Nutrition 21, LLC



Directions: To assess tolerance, take (1) capsule on an empty stomach 30-45 minutes before breakfast and an additional (1) capsule 5-6 hours later. Once tolerance has been assessed, an additional (1) capsule may be added to the morning serving. Under no circumstances should the initial serving be exceeded or the warnings on this bottle be ignored. **DO NOT EXCEED 3 CAPSULES IN ANY 24 HOUR PERIOD. DO NOT USE PRODUCT FOR LONGER THAN 8 WEEKS FOLLOWED BY A SUBSEQUENT 4 WEEK BREAK.**

WARNING: DO NOT USE IN COMBINATION WITH CAFFEINE OR ANY STIMULANTS FROM OTHER SOURCES INCLUDING BUT NOT LIMITED TO COFFEE, TEA, SODA, AND OTHER DIETARY SUPPLEMENTS OR MEDICATIONS. DO NOT COMBINE WITH ALCOHOL. Do not use under extreme conditions of heat, sleep deprivation, or dehydration. This product is only intended to be consumed by healthy adults 18 years of age or older. Consult with your physician before using this product, especially if you are using any prescription or over the counter medication or if you have pre-existing medical condition including but not limited to: high or low blood pressure, cardiac arrhythmia, stroke, heart, liver, kidney, or thyroid disease, seizure disorder, psychiatric disease, diabetes, difficulty urinating due to prostate enlargement or if you are taking a MAOI (Monoamine Oxidase Inhibitor) or any other medication. Do not take this product if you are pregnant or nursing. Discontinue use 2 weeks prior to surgery. Discontinue use and consult your health care professional if you experience an adverse reaction to this product. Do not exceed recommended serving. Do not use if safety seal is broken or missing. **KEEP OUT OF REACH OF CHILDREN.**

†These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.