This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose. treat, cure, or prevent any disease.


KEEP OUT OF REACH OF CHILDREN Store in a cool, dry place,

For More Information: 1-888-462-2548 SHOP NOW @ GNC.COM

Distributed by: **General Nutrition Corporation** Pittsburgh, PA 15222 USA



## GNC milestones<sup>t</sup>



No Dairy 120 CAPLETS CODE 200812

Serving Size Two Caplets

Servings Per Container 60

(as Ascorbic Acid & Calcium Ascorbate)

Amount Per Serving

Directions: As a dietary supplement, take two caplets daily with food

(400 mcg Folic Acid)

## Amount Per Serving % Daily Value Supplement Facts Phosphorus (as Dicalcium Phosphate) 50 mg 4% lodine (as Potassium Iodide) 150 mcg 100% % Daily Value Zinc (as Zinc Oxide) \_\_\_\_\_\_\_15 mg\_\_\_\_136% Conner (as Cunric Sulfate) 2 mg 222% Manganese (as Manganese Sulfate). 2 mg. 87% Vitamin E (as d-alpha Tocophery Succinate) ..... 24,79 mg ..... 165% Chloride (as Potassium Chloride) 47 mg 2% Vitamin K (as Phytonadione) 80 mcg 67% Thiamin (Vitamin B-1) (as Thiamin Mononitrate) .... 4.5 mg.... 375% Riboflavin (Vitamin B-2) \_\_\_\_\_\_\_ 5,1 mg\_\_\_ 392% Cranberry Fruit Concentrate 80 mg Miscin /as Miscingmide) 20 mg 125% (Vaccinium macrocarporti Fotate \_\_\_\_\_\_\_667 mcg DFE ... 167% Flax Seed Hulls (Linum usitatissimum) 50 mg (Vaccinium corymbosum) Lycopene 500 mcg 4 Zeaxanthin......200 mcg.......

DSG

OTHER INGREDIENTS: Cellulose, Stearic Acid Vegetable Source, Croscarmellose Sodium, Silicon Dioxide Magnesium Stearate Vegetable Source, Titanium Dioxide (Mineral Whitener), Vegetable Acetoglycerides, Polyethylene Glycol, Polysorbate 80, Caramel Color, Ethyl Vanillin,

\* Daily Value not petablished

CONTAINS: Soybeans and Wheat.

Phosphate, Calcium Ascorbate, Calcium D-Pantothenate)

WARNING: Consult your physician prior to using this product if you are pregnant, nursing, taking medication, or have a medical condition. Discontinue use two weeks prior to surgery.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.