Garcinia cambogia is a type of fruit native to Indonesia, Asia, and Africa. Traditionally, it has been used to make different foods such as curry and soup and to make meals more filling. An active constituent of the fruit, HCA (hydroxycitric acid), is believed to have beneficial effects on appetite and weight management.



Warnings: Not intended for use by persons under age 18. Do not exceed recommended dose. Do not take for more than eight (8) consecutive weeks. This product should not be taken by pregnant or lactating women. Not intended for use by those with a medical condition. Get the consent of a licensed physician before using this product, especially if you are taking medication, have a medical condition. or thinking about becoming pregnant. Discontinue use two weeks prior to surgery. KEEP THIS PRODUCT AND ALL SUPPLEMENTS OUT OF THE REACH OF CHILDREN, DO NOT USE IF **SAFETY SEAL IS BROKEN** 

- Store at 15-30°C (59-86°F)
- Protect from heat, light and moisture.
- Do not purchase if seal is broken.

\* THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE. TREAT, CURE, OR PREVENT ANY DISEASE.

<sup>1</sup>When combined with a proper exercise and nutrition regimen. Statements based on early-stage independent 3rd party in vivo and / or in vitro model scientific research data findings for individual ingredients.



## GARGINIA ONCENTRATED

1500mg PURE GARCINIA CAMBOGIA EXTRACT 50.0% Hydroxycitric Acid (HCA)

**MAY EXPERIENCE:** 

WEIGHT LOSS\*+

APPETITE SUPPORT\*†





**DIETARY SUPPLEMENT** 

Please read entire label before use.

## SUPPLEMENT FACTS

Serving Size 1 Tablet Servings Per Container 60

## **Amount Per Serving**

% Daily Value

Garcinia cambogia fruit extract

1.5 g \*

(50% hydroxycitric acid)

\*\* Daily Value not established.

Other Ingredients: Microcrystalline cellulose, calcium carbonate, coating (hypromellose, hydroxypropyl cellulose, and polyethylene glycol), croscarmellose sodium, stearic acid, magnesium stearate, and silica.

**Suggested Use:** Take one (1) tablet after exercise or 30-45 minutes before the heaviest meal of the day, or as directed by a qualified healthcare practitioner.



Manufactured for and Distributed By: BPI Health. 3149 SW 42nd St. Suite 200 Hollywood, FI, 33312

To report an adverse event or for more information call: 954,926,0900 (tel).