NATURAL TESTOSTERONE BOOSTER'

DAA TEST-5 IS DESIGNED TO BOOST TESTOSTERONE LEVELS

BY ACTIVATING

PATHWAYS

IN THE BODY. THIS RELEASES

NATURAL TESTOSTERONE

TO SUPPORT INCREASED

STRENGTH, ENERGY, MUSCLE MASS, RECOVERY AND LIBIDO'





DIETARY SUPPLEMENT 96 CAPSULES

STRENGTH[†]

LIBIDO[†]





SUGGESTED USE: As a dietary supplement, adults take 1 serving (4 capsules) once a day. Recommended dosing regimen includes taking one serving for twelve straight days, followed by two days off before repeating. Please read the entire label before use. Should not be consumed for more than 4 consecutive weeks with a 2 to 4 week washout period between cycles.

Supplement Facts

Serving Size: 4 Capsules Servings Per Container: 24

	Amount Per Serving	% Daily Value
DAA Test 5 Complex [†]	3,296 mg	**
Sodium D-Aspartic Acid	3,120 mg	**
Testosterone Support Complex	t 176 mg	**
Chrysin, Fenugreek (<i>Trigonella foenum-graecum</i>) extract (seed) (std. 80% Glycosides), <i>Eurycoma lonigifolia</i> extract (root) 20:1, Diindolylmethane (DIM)		

** Daily Value not established.

Other Ingredients: Microcrystalline Cellulose, Gelatin, Magnesium Stearate, Titanium Dioxide, Silicon Dioxide

This product was produced in a facility that may also process ingredients containing milk, eggs, fish, shellfish, tree nuts, peanuts, wheat, and soybeans.

KSM-66® is a registered trademark of Ixoreal Biomed Private Limited.

WARNING: DO NOT USE IF PREGNANT, NURSING, OR PLAN ON BECOMING PREGNANT. THIS PRODUCT IS INTENDED FOR USE BY HEALTHY ADULTS OVER THE AGE OF 18. Do not use this

PROUDD 15 INTENDED FOR USE BY INSLIFY ADDUS OVER INF. ARE OF 18. Di Ant Use this product continuosily for more than 8 weeks. Do not comhine with alcohol. Not intended for use by those with a medical condition. Use only as directed. Do not exceed recommended daily intake. Consult a physician or licensed qualified health care professional profit to use if you are pregnant or nursing, or if you are taking medication, including, but not limited to MAD inhibitors, antidepressants, aspirin, monsteroidal anti-inflammatory drugs or products containing phenylephrine, ephedrine, pseudoephedrine, or other stimulants, or if you have a medical condition, including, but not limited to heart. Jiver, kidney, or thyroid disease, psychiatric or epileptic disorders, difficulty unintang, diabetes, high blood ressure, cardiae arrhythmia, recurrent headches, enlarged prostate or glaucoma. Discontinue 2 weeks prior to surgery or if you experience rapid heartbeat, dizziness, severe headache or shortness of breath. DD NOT USE IF SAFETY SEU. UNDER CAP IS BROKEN. KEEP OUT G FREACH OF CHIDREN.

Store in a cool, dry place. Please recycle.

¹These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.