

**GNC**  
TOTAL LEAN®

TRIPLE STRENGTH  
**L-CARNITINE**  
**LIQUID**

DIETARY SUPPLEMENT



3 g of L-Carnitine per serving, ultra-concentrated

Supports fatty acid metabolism &  
helps to fuel muscle recovery\*

Sugar Free

**FRUIT PUNCH**

NATURAL & ARTIFICIAL FLAVORS



16 FL.OZ (1 PINT) (473 ML)

ABOUT 30 SERVINGS

**OUR QUALITY  
COMMITMENT TO YOU**

Since 1935, our mission as the industry leader has been to provide you with the highest-quality nutritional supplements. We continue to stand behind this mission. If you're not 100% satisfied with your purchase, return the unused portion of the product with your receipt within 30 days and we'll either refund your money or find a replacement product that is right for you.

**KEEP OUT OF REACH OF  
CHILDREN.**

Refrigerate after opening.

For More Information:

1-888-462-2548

GNC.com

Distributed by:

General Nutrition Corporation

Pittsburgh, PA 15222 USA

K1064804



**CODE 732281**

**GTG**

**DIRECTIONS:** As a dietary supplement, take one tablespoon daily. May be taken directly or added to water or your favorite beverage. For best results, take 30-60 minutes prior to exercise. Shake well before each use. Refrigerate after opening.

## Supplement Facts

Serving Size One Tablespoon (15 mL)

Servings Per Container About 30

Amount Per Serving		% Daily Value
Calories	20	
Total Carbohydrate	0 g	0% †
Total Sugars	0 g	*
Includes 0 g Added Sugars		0% †
Vitamin B-6 (as Pyridoxine HCl)	2 mg	118%
Pantothenic Acid (as D-calcium Pantothenate)	25 mg	500%
L-Carnitine (as L-Carnitine and L-Carnitine HCl)	3 g	*

† Percent Daily Values are based on a 2,000 calorie diet.

\* Daily Value not established.

**OTHER INGREDIENTS:** Purified Water, Vegetable Glycerin, Citric Acid, Natural and Artificial Flavors, Potassium Sorbate (Preservative), Sodium Hexametaphosphate (Preservative), Calcium Propionate (Preservative), Sucralose, Acesulfame Potassium and FD&C Red #40.

**WARNING:** Consult your physician prior to using this product if you are pregnant, nursing, taking medication, or have a medical condition. Discontinue use two weeks prior to surgery.

\* This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.