THE OBJECTIVE:

WHEN IT'S TIME TO GO TO BATTLE, YOU GO HARD WITH TOTAL War. Experience a workout with intensity you didn't Think was possible. New found strength and energy is Just Minutes away. Lock-in and get ready to bring the noise!*

TOTAL WAR SUGGESTED USE: WHEN IT COMES TIME TO BRING THE NOISE FOR A WORKOUT, ASSESS YOUR TOLERANCE BY TAKING ½ SCOOP PRIOR TO TRAINING. ONCE YOUR TOLERANCE IS ASSESSED MIX (1) SCOOP WITH 4-6 OUNCES 30 MINUTES PRIOR TO TRAINING. DO NOT EXCEED MORE THAN (1) SCOOP PER DAY.



			σ
Supplement Serving Size: 1 Scoop (14.7g)	Fac	cts	45
Servings Per Container: 30			191
Amount Per	Serving	% DV	
L-Citrulline DL-Malate 2:1 Beta-Alanine	6g 3.2g	**	754
Agmatine Sulfate Taurine	1g 1g	** **	
Caffeine Anhydrous AMPiblast™ (Juniperus Communis)(berry)	250mg 150ma	** **	
Di-Caffeine Malate Cocophenols ™ (Theobroma cacao & Camelia sinensis Extract)	100 mg	** **	not asse.
Naringin 98% (Citrus Grandis) (Fruit) Bioperine™ (Black Pepper Extract)(Fruit)	25mg 10mg	**	dise is by
*Percent Daily Values are based on a 2000 colorie diet **Daily Value not established			an evaluated This product prevent any
heringesien: Naturi end erfähilferer, Desdam phosphete, Sala, Galaan skaaet, Sanatova, GA NANUFACTURED FOR AND DIS 1. EDCONI SUPPLEMENTS 1. 41 SROGERSCIRCLE SUITE 3BOCA	nicocid, Mohodentria, FRIBUT L: 561-9 RATON,	Analone patroion. 5 E D B Y : 9 6 1 - 8 3 4 9 F L 3 3 4 8 7	nts have not been evaluated Administration. This product ose, treat, cure, or prevent any
VARNING, KEP OUT OF THE REACH OF CHILDREN. be consumed by healthy dubit. If years of ages of consumed and the second provide a second provide and inder to hale or two blood presence collaborations and the second provide and the second provide and the second provide a second provide a second provide the second provide and the second provide and the second provide a second provide a second to a second provide a second provide a second to second provide a second provide a second provide to second provide a second provide a second provide to second provide a second provide a second provide a second research provide a second provide a second provide a second research provide a second provide a second provide a second research provide a second provide a second provide a second research provide a second provide a second provide a second research provide a second provide a second provide a second research provide a second provide a second provide a second research provide a second provide a second provide a second research provide a second provide a second provide a second research provide a second provide a second provide a second provide a second research provide a second provide	r older.Befc Avoid usin idition inclu nia, stroke, h niatric disea if you are prine to de consult you	re using this githis product ding but not seart, liver or se, diabetes, teking MAO 2 weeks prior shydration or r healthcare RE. PRODUCT	 These statements he Food and Drug Admin intended to diagnose, tr

RED